

Quantum Decoherence and Higher Order Corrections to the Large Time Exponential Behaviour

I.Ya. Aref'eva* and I.V. Volovich [†]

*Steklov Mathematical Institute, Russian Academy of Sciences
Gubkin St.8, GSP-1, 117966, Moscow, Russia*

Abstract

There exists the well known approximate expression describing the large time behaviour of matrix elements of the evolution operator in quantum theory: $\langle U(t) \rangle \simeq \exp(at)$. This expression plays the crucial role in considerations of problems of quantum decoherence, radiation, decay, scattering theory, stochastic limit, derivation of master and kinetic equations etc. This expression was obtained in the Weisskopf-Wigner approximation and in the van Hove (stochastic) limit. We derive the exact general formula which includes the higher order corrections to the above approximate expression: $\langle U(t) \rangle = \exp(At + B + C(t))$. The constants A and B and the oscillating function $C(t)$ are computed in perturbation theory. The method of perturbation of spectra and renormalized wave operators is used. The formula is valid for a general class of Hamiltonians used in statistical physics and quantum field theory.

*arefeva@mi.ras.ru

[†]volovich@mi.ras.ru

1 Introduction

The study of the large time behaviour of the evolution operator in statistical physics and quantum field theory is the subject of numerous investigations. The basic object to study in quantum field theory is the scattering matrix. The physical idea behind the scattering matrix approach is that in the scattering processes there exists a characteristic time scale such that in a time regime larger than this time scale one can neglect interaction and particles evolve according to the free dynamics [1]. However, due to the infinite number of degrees of freedom in quantum field theory asymptotic dynamics is not simply governed by the free Hamiltonian. There are effects of renormalization of vacuum energy and one particle states and the asymptotic states become in fact the states of dressed particles [2]. Therefore one has to deal with renormalized, or dressed wave operators [3, 4, 5, 6]. In this paper we will use the method of renormalized wave operators to compute matrix elements of the evolution operator for finite time.

There are many important problems in quantum field theory where we are interested in the large but not infinite time and where the standard S -matrix description is not very convenient or even not applicable. These include processes with unstable particles [7, 8] (in fact almost all particles are unstable), atom-photon interactions [9], elementary particles in "semidressed states" with non-equilibrium proper fields [10], electroweak baryogenesis and phase transitions in the early Universe and in high-energy collisions [11], quantum optics [12], quantum decoherence (see for example [12, 13, 14]) etc. In the consideration of such processes we are interested in the time regime smaller than the "infinite" time when the S -matrix description becomes applicable. The consideration of such processes belongs to non-equilibrium quantum field theory, see [15] for more discussions.

In statistical physics there is not just one but several relevant time scales and as a result we don't have here a universal method comparable with the S -matrix approach in quantum field theory. One can say that the role of S -matrix approach in non-equilibrium statistical physics is played by various master and kinetic equations. Various methods of consideration of time evolution for classical and quantum systems have been developed by Bogoliubov [16], Weisskopf and Wigner (see [8]), van Hove [18], Prigogine [19] and many others, for a review see for example [20, 21].

A general method in non-equilibrium statistical physics and quantum field theory is the method of stochastic limit, see [22, 23]. The idea of this method is the systematic application of the $\lambda^2 t$ -limit and quantum stochastic differential equations. One considers the evolution operator $U(t)$ of quantum system for small coupling constant λ and large time t . The limit

$$\lambda \rightarrow 0, \quad t \rightarrow \infty, \quad \lambda^2 t = \text{fixed} = \tau \quad (1.1)$$

has been considered by Bogoliubov [16], Friedrichs [17] and van Hove [18]. In the quantum theory of open systems, the limit (1.1) is known as the van Hove or the $\lambda^2 t$ limit.

In this paper we study corrections to the (van Hove) stochastic limit. For this purpose a general formula for the matrix elements of the evolution operator is obtained. This formula is used for the investigation of the large time and small coupling constant asymptotic behaviour. We consider a very general class of Hamiltonians used in solid state physics and quantum field theory

$$H = H_0 + \lambda V, \quad (1.2)$$

where H_0 is a free Hamiltonian, V describes an interaction and λ is the coupling constant.

It is well known that the large time behaviour of the vacuum expectation value of the evolution operator

$$U(t) = e^{itH_0} e^{-itH}, \quad (1.3)$$

is given by

$$\langle U(t) \rangle \simeq e^{at}, \quad (1.4)$$

where a is a constant. This expression can be obtained in the Weisskopf-Wigner approximation [8] or in the van Hove (stochastic) limit. In this paper we obtain the following exact general formula valid for any time t

$$\langle U(t) \rangle = e^{iAt+B+C(t)}. \quad (1.5)$$

Here A and B are constants for which a representation in perturbation theory will be given and $C(t)$ is a function which under rather general assumptions can be represented for large time t as

$$C(t) = \frac{f(t)}{t^\alpha}. \quad (1.6)$$

Here $f(t)$ is a bounded oscillating function and the exponent α depends on the model and on the dimension of space ($\alpha = 3/2$ for the physical 3-dimensional space and for a general class of models). We derive the main formula (1.5) by using the theory of perturbation of spectra and renormalized wave operators.

One of the remarkable features of quantum mechanics which most distinguishes it from classical mechanics is the coherent superposition of physical states. The important point is the physical distinction between the coherent superposition of states and the classical mixture, see for example [12, 13]. In particular the maintenance of quantum coherence is a crucial requirement of the ability of quantum computers to be more efficient in certain problems than classical computers (see for example [14] and refs therein). Any state of a quantum system can be described by the off-diagonal elements of the density operator. In fact the dominant contribution for large time to any matrix element of the evolution operator comes from vacuum and if $\Re a < 0$ in (1.4) then one has the exponential decay. The constant a is a function of the coupling constant and other parameters of the model. To suppress decoherence we would like to have the regime with $\Re a = 0$. In such a case we have to investigate corrections to the approximate expression (1.4). The solution of this problem is given in this paper and it is presented in (1.5).

Expectation value in (1.5) is taken over vacuum. For the case of one-particle states we obtain

$$\langle p|U(t)|p' \rangle = e^{iA(p)t+B(p)+C(t,p)} \delta(p - p'). \quad (1.7)$$

The formulae (1.5) and (1.7) have a very general character. We prove (1.5) in Section 3 for a very general class of Hamiltonians. From this formula one gets the stochastic limit of evolution operator

$$\langle U(\tau/\lambda^2) \rangle = e^{iA_2\tau}(1 + o(\lambda)), \quad (1.8)$$

as well as corrections to stochastic limit

$$\langle U(\tau/\lambda^2) \rangle = e^{iA_2\tau + \lambda^2(B_2 + i\tau A_4) + o(\lambda^2)}. \quad (1.9)$$

In Section 4 we calculate corrections to the stochastic limit for the one particle matrix elements of the evolution operator for the case of translation invariant Hamiltonians

$$\langle p|U(\tau/\lambda^2)|p'\rangle = e^{iA_2(p)\tau}(1 + \lambda^2(B_2(p) + i\tau A_4(p)) + o(\lambda^2))\delta(p - p') \quad (1.10)$$

The class of considered Hamiltonians includes the Bose and Fermi gases, phonon self-interaction and electron-phonon interaction, quantum electrodynamics in external fields etc.

We give two independent derivations of formula (1.5) and (1.10). The first method consists in the direct examination of perturbation theory. The second one uses powerful results of spectral theory and renormalized wave operators [6]. Although the second method is simpler we present both of them since the first one can be used also in the case of decay when the results of the standard scattering theory are not applicable.

2 Notations and auxiliary results

2.1 Hamiltonians

We consider Hamiltonians of the form (1.2) where H_0 is a free Hamiltonian

$$H_0 = \sum_i \int \omega_i(k) a_i^*(k) a_i(k) d^d k \quad (2.1)$$

and V is the sum of Wick monomials. Creation and annihilation operators $a_i^*(k)$, $a_i(k)$ describe particles or quasiparticles and they satisfy the commutation or anticommutation relations

$$[a_i(k), a_j^*(k')]_{\pm} = \delta_{ij} \delta(k - k') \quad (2.2)$$

Here $k, k' \in R^d$ and $i, j = 1, \dots, N$ label the finite number of different types of (quasi)particles. Examples of one-particle energy $\omega_i(k)$ include the relativistic ($\omega(k) = (k^2 + m^2)^{1/2}$) and non-relativistic ($\omega(k) = k^2/2 - \omega_0$) laws, the Bogoliubov spectrum ($\omega(k) = (bk^4 + k^2 v(k))^{1/2}$), the Fermi quasiparticle spectrum ($\omega(k) = |k^2/2m - \mu|$) etc.

We consider two different types of Wick polynomials. The first type describes an interaction in the case when there is not the translation invariance

$$V = \sum_{I,J} \int v(p_1, i_1 \dots p_I, i_I | q_1, j_1 \dots q_J, j_J) \prod_{l=1}^I a_{i_l}^*(p_l) dp_l \prod_{r=1}^J a_{j_r}(q_r) dq_r \quad (2.3)$$

were $v(p_1 \dots p_I | q_1 \dots q_J)$ are some test functions.

The second type is described by the translation invariant Hamiltonian

$$V = \sum_{I,J} V_{I,J} = \sum_{I,J} \int \hat{v}(p_1, i_1, \dots, p_I, i_I | q_1, j_1 \dots q_{J-1}, j_{J-1}, J_J) \delta \left(\sum_l^I p_l - \sum_r^J q_r \right) \prod_{l=1}^I a_{i_l}^*(p_l) dp_l \prod_{r=1}^J a_{j_r}(q_r) dq_r \quad (2.4)$$

Clearly the delta function causes the trouble and there are singular terms in (2.4). Namely, $V_{I,0}\phi_n$ does not belong to the Fock space unless $\phi_n = 0$. This singularity is called the volume singularity. To give a meaning to the Hamiltonian with interaction (2.4) one has to introduce a volume cut-off, then perform the vacuum renormalization and vacuum dressing and only after that remove the cut-off. This procedure defines the hamiltonian in a new space (see [5, 24] for details). To avoid this difficulty in this paper we will assume that for translation invariant interaction there are no pure creation and annihilation terms.

2.2 Evolution operator

We will investigate the evolution operator

$$U(t) = e^{itH_0} e^{-it(H_0 + \lambda V)}. \quad (2.5)$$

In perturbation theory the evolution operator (2.5) has the representation

$$U(t) = 1 - i\lambda \int_0^t V(t_1) dt_1 + (-i\lambda)^2 \int_0^t dt_1 \int_0^{t_1} dt_2 V(t_1) V(t_2) + \dots \quad (2.6)$$

where

$$V(t) = e^{itH_0} V e^{-itH_0}$$

We will also use the evolution operator with the adiabatic cut-off

$$U_\epsilon(t) = 1 - i\lambda \int_0^t V_\epsilon(t) U_\epsilon(t) dt \quad (2.7)$$

$$V_\epsilon(t) = e^{-\epsilon|t|} e^{itH_0} V e^{-itH_0}$$

In this case one also has the perturbation series

$$U_\epsilon(t) = 1 - i\lambda \int_0^t V_\epsilon(t) dt + (-i\lambda)^2 \int_0^t dt_1 \int_0^{t_1} dt_2 V_\epsilon(t_1) V_\epsilon(t_2) + \dots \quad (2.8)$$

We will use for $V_{I,J}$ the diagram representation. The corresponding diagram has one vertex and I lines going from the vertex to the left and J lines going to the right. The first I lines represent creation operators and the last J lines represent annihilation operators. In what follows we will use the Wick theorem:

$$V_{I,J} W_{N,M} =: V_{I,J} W_{N,M} : + \sum_s^{\min\{J,N\}} V_{I,J} \underbrace{\text{---} \circ \text{---}}_s W_{N,M}. \quad (2.9)$$

The kernel of the Wick monomial $: V_{I,J} W_{N,M} :$ is

$$v_{I,J} \otimes w_{N,M} \quad (2.10)$$

and $V_{I,J} \underbrace{\text{---} \circ \text{---}}_s W_{N,M}$ is the Wick monomial

$$V_{I,J} \underbrace{- \circ -}_s W_{N,M} = \int \prod_{r=1}^{I+N-s} (dp_r a^*(p_r)) \cdot \prod_{l=1}^{J+M-s} (dq_l a(q_l)) C_J^s C_N^s t! \int \prod_{r=1}^s dk_r (v \circ_s w)(k_1, \dots, k_r, p_1, \dots, q_1 \dots) \quad (2.11)$$

with the following kernel

$$(v \circ_s w)(k_1, \dots, k_r, p_1, \dots, q_1 \dots) = \quad (2.12)$$

$$v_{I,J}(p_1, \dots, p_I; k_1, \dots, k_s, q_1, \dots, q_{J-s}) w_{N,M}(k_1, \dots, k_s; p_{I+1}, \dots, p_{I+N-1}, q_{J-s+1}, \dots, q_{J-s+M}).$$

Below we will drop out the symbol s in $\circ_s$ and to specify the concrete form of contractions will refer to corresponding diagrams.

The equality (2.9) for the 4-point interaction has the diagram representation as shown on Fig. 1.




Figure 1: Graphs representation of (2.9)

We will use also the following notations. The line of the graph is called the internal if it connects two vertices of the graph. A graph is the connected graph if all its vertices are connected by a set of internal lines otherwise it is called the disconnected one. A connected graph is called the one-particle reducible (1PR) if after the removing a line it becomes disconnected. A connected graph is called the one-particle irreducible (1PI) if after the removing any line it is still connected.

The following "linked cluster theorem" [5] will be used:

$$U(t) =: e^{U_c(t)} : \quad (2.13)$$

where

$$U_c(t) = \sum_{n=1}^{\infty} (-i\lambda)^n \int_0^t dt_1 \dots \int_0^{t_{n-1}} dt_n (V(t_1) \dots V(t_n))_c$$

Here the index c in U_c indicates that one takes only the connected diagrams.

The similar relation is true for evolution operator with the adiabatic cutoff.

Below for simplicity of notations we consider interactions with the only one type of particles, but the main results are valid for arbitrary number of types of particles.

3 Evolution operator for non-translation invariant Hamiltonians

3.1 Second order

For the vacuum matrix element of the evolution operator we obtain from (2.13) the representation:

$$\langle 0|U(t)|0\rangle = e^{\mathcal{E}(t)} \quad (3.1)$$

where

$$\mathcal{E}(t) = \langle 0|(-i\lambda \int_0^t dt_1 V(t) + (-i\lambda)^2 \int_0^t dt_1 \int_0^{t_1} dt_2 V(t_1)V(t_2) + \dots)_c|0\rangle. \quad (3.2)$$

Here the symbol $(\dots)_c$ means that we keep only connected diagrams.

Representation (3.1), (3.2) permits us to calculate the leading terms of the asymptotic behaviour of the matrix elements of the evolution operator for large time t as well the corrections to the leading terms. In fact we will show that $\mathcal{E}(t)$ has the following form

$$\mathcal{E}(t) = At + B + C(t) \quad (3.3)$$

where one has the perturbative expansions

$$A = \lambda^2 A_2 + \lambda^3 A_3 + \dots, \quad B = \lambda^2 B_2 + \lambda^3 B_3 + \dots, \quad C(t) = \lambda^2 C_2(t) + \lambda^3 C_3(t) + \dots \quad (3.4)$$

and $C_n(t)$ vanishes for large t .

Let us find explicitly these terms in the second order of perturbation theory for the Hamiltonian

$$H = H_0 + \lambda V, \quad (3.5)$$

where

$$H_0 = \int \omega(p) a^*(p) a(p) dp \quad (3.6)$$

and the interaction has the form

$$V = \int (v(p_1, \dots, p_n) a^*(p_1) \dots a^*(p_n) + c.c.) dp_1 \dots dp_n \quad (3.7)$$

Here $\omega(p)$ is a positive smooth function, for example $\omega(p) = \sqrt{p^2 + m^2}$, $m > 0$ and $v(p_1, \dots, p_n)$ is a test function. For this interaction the first term in (3.2) is identically zero. The second term in (3.2) equals to

$$\mathcal{E}^{(2)}(t) = (-i\lambda)^2 \int dp_1 \dots dp_n |v(p_1, \dots, p_n)|^2 \int_0^t dt_1 \int_0^{t_1} dt_2 e^{it_1 E_1 + it_2 E_2} \quad (3.8)$$

where

$$E_2 = -E_1 = E(p_1, \dots, p_n) = \sum_{i=1}^n \omega(p_i) \quad (3.9)$$

By using the equality

$$\int_0^t dt_1 \int_0^{t_1} dt_2 e^{-it_1 E + it_2 E} = -\frac{i}{E}t + \frac{1}{E^2} - \frac{1}{E^2}e^{-itE} \quad (3.10)$$

we get

$$\mathcal{E}^{(2)}(t) = (-i\lambda)^2 \int dp_1 \dots dp_n |v(p_1, \dots, p_n)|^2 \left(-\frac{i}{E}t + \frac{1}{E^2} - \frac{1}{E^2}e^{-itE}\right) \quad (3.11)$$

Therefore we obtain the expression of the form (3.3)

$$\mathcal{E}(t) = \lambda^2 A_2 t + \lambda^2 B_2 + \lambda^2 C_2(t) + \dots \quad (3.12)$$

where

$$A_2 = i \int \frac{|v(p_1, \dots, p_n)|^2}{E(p_1, \dots, p_n)} dp_1 \dots dp_n, \quad (3.13)$$

$$B_2 = - \int \frac{|v(p_1, \dots, p_n)|^2}{E(p_1, \dots, p_n)^2} dp_1 \dots dp_n, \quad (3.14)$$

$$C_2(t) = \int \frac{|v(p_1, \dots, p_n)|^2}{E(p_1, \dots, p_n)^2} e^{-itE(p_1, \dots, p_n)} dp_1 \dots dp_n \quad (3.15)$$

We have obtained the following

Theorem 1. The vacuum expectation value of the evolution operator for the Hamiltonian (3.5) in the second order of perturbation theory has the form

$$\langle U(t) \rangle = e^{\lambda^2 A_2 t + \lambda^2 B_2 + \lambda^2 C_2(t)} \quad (3.16)$$

where A_2 , B_2 and $C_2(t)$ are given by (3.13), (3.14) and (3.15).

Remark. By using the stationary phase method one can prove that the function $C_2(t)$ vanishes as $t \rightarrow \infty$ (see below).

3.2 Decay

We have proved theorem 1 under the assumption $\omega(p) > 0$. However the obtained formula (3.16) is valid in the more general case when one has the decay. In this case formula (3.16) still is true but in the expressions (3.13)-(3.15) one has to substitute $E \rightarrow E - i0$. Let us consider the important case when

$$\omega(p) = \frac{p^2}{2} - \omega_0, \quad \omega_0 > 0 \quad (3.17)$$

Instead of (3.10) we will use now the identity

$$\int_0^t dt_1 \int_0^{t_1} dt_2 e^{i(t_2 - t_1)E} = t \int_0^t \left(1 - \frac{\sigma}{t}\right) e^{-i\sigma E} d\sigma. \quad (3.18)$$

We have

$$\mathcal{E}^{(2)}(t) = (-i\lambda)^2 \int dp_1 \dots dp_n |v(p_1, \dots, p_n)|^2 \int_0^t dt_1 \int_0^{t_1} dt_2 e^{i(t_2 - t_1)E(p_1, \dots, p_n)} \quad (3.19)$$

$$\begin{aligned}
&= -\lambda^2 t \int_0^t d\sigma \left(1 - \frac{\sigma}{t}\right) \int dp_1 \dots dp_n |v(p_1, \dots, p_n)|^2 e^{-i\sigma E(p_1, \dots, p_n)} \\
&= \lambda^2 t A_2(t) + \lambda^2 B_2(t)
\end{aligned}$$

where

$$A_2(t) = - \int_0^t d\sigma F(\sigma), \quad B_2(t) = \int_0^t d\sigma \sigma F(\sigma) \quad (3.20)$$

and

$$F(\sigma) = \int dp_1 \dots dp_n |v(p_1, \dots, p_n)|^2 e^{-i\sigma E(p_1, \dots, p_n)}. \quad (3.21)$$

By using the stationary phase method we obtain the following asymptotic behaviour of the function $F(\sigma)$ as $\sigma \rightarrow \infty$:

$$F(\sigma) = \left(\frac{2\pi i}{\sigma}\right)^{dn/2} e^{i\sigma\omega_0} |v(0)|^2 \left[1 + o\left(\frac{1}{\sigma}\right)\right].$$

Therefore for $dn \geq 3$ there exist the limits

$$\lim_{t \rightarrow \infty} A_2(t) = A_2 = - \int_0^\infty d\sigma F(\sigma) \quad (3.22)$$

$$\lim_{t \rightarrow \infty} B_2(t) = B_2 = \int_0^\infty d\sigma \sigma F(\sigma)$$

because there exists the limit

$$\lim_{t \rightarrow \infty} \int_1^t e^{i\sigma\omega_0} \frac{d\sigma}{\sigma^{1/2}}.$$

Moreover one has

$$A_2(t) = - \int_0^\infty d\sigma F(\sigma) + o\left(\frac{1}{t^2}\right), \quad B_2(t) = \int_0^\infty d\sigma \sigma F(\sigma) + o\left(\frac{1}{t}\right).$$

If $dn \geq 5$ one gets also

$$A_2 = i \int dp_1 \dots dp_n \frac{|v(p_1, \dots, p_n)|^2}{E(p_1, \dots, p_n) - i0} \quad (3.23)$$

$$B_2 = - \int dp_1 \dots dp_n \frac{|v(p_1, \dots, p_n)|^2}{(E(p_1, \dots, p_n) - i0)^2}. \quad (3.24)$$

Indeed one has

$$A_2 = - \int_0^\infty d\sigma F(\sigma) = - \lim_{\epsilon \rightarrow 0} \int_0^\infty d\sigma F(\sigma) e^{-\sigma\epsilon}.$$

This is true due to the Lebesgue theorem since $|F(\sigma)e^{-\sigma\epsilon}| \leq |F(\sigma)|$ and $F(\sigma) \in L_1(R_+)$ (L_1 is the space of absolute integrable functions) if $nd \geq 3$. Substituting in the above formula the representation (3.21) and changing the order of integrations (we can do this due to the Fubini theorem since for positive ϵ the function $|v(p_1, \dots, p_n)|^2 e^{-i\sigma(E(p_1, \dots, p_n) - i\epsilon)}$ belongs to the space $L_1(R_+ \times R^{nd})$ of absolute integrable functions), we can perform the integration over σ explicitly

$$A_2 = \lim_{\epsilon \rightarrow 0} (i) \int dp_1 \dots dp_n \frac{|v(p_1, \dots, p_n)|^2}{E(p_1, \dots, p_n) - i\epsilon} = i \int dp_1 \dots dp_n \frac{|v(p_1, \dots, p_n)|^2}{E(p_1, \dots, p_n) - i0}.$$

The same calculation is true for B_2 with the more strong assumption : $dn \geq 5$,

$$\begin{aligned} B_2 &= \int_0^\infty d\sigma \sigma F(\sigma) = \lim_{\epsilon \rightarrow 0} \int_0^\infty d\sigma \sigma \int dp_1 \dots dp_n |v(p_1, \dots, p_n)|^2 e^{-i\sigma(E(p_1, \dots, p_n) - i\epsilon)} \\ &= -\lim_{\epsilon \rightarrow 0} \int dp_1 \dots dp_n \frac{|v(p_1, \dots, p_n)|^2}{(E(p_1, \dots, p_n) - i\epsilon)^2} = -\int dp_1 \dots dp_n \frac{|v(p_1, \dots, p_n)|^2}{(E(p_1, \dots, p_n) - i0)^2}. \end{aligned}$$

We have proved the following theorem.

Theorem 2. The asymptotic behaviour as $t \rightarrow \infty$ of the vacuum expectation value of the evolution operator for the Hamiltonian (3.5) with the dispersion law (3.17) in the second order of perturbation theory is

$$\langle U(t) \rangle = e^{\lambda^2 A_2 t + \lambda^2 B_2 + \lambda^2 o(1/t)} \quad (3.25)$$

where A_2 and B_2 are given by (3.22) (or (3.23) and (3.24)). After the rescaling $t \rightarrow t/\lambda^2$ one gets the λ^2 corrections to the stochastic limit

$$\langle U(t/\lambda^2) \rangle = e^{A_2 t + \lambda^2 B_2 + \lambda^2 o(\lambda^2/t)}. \quad (3.26)$$

3.3 Example

We discuss here the evolution operator for the simple explicitly solvable model described by the Hamiltonian

$$H = \int \omega(k) a^+(k) a(k) d^d k + \lambda \int (a(k) \bar{v}(k) + a^*(k) v(k)) d^d k. \quad (3.27)$$

We will see that the vacuum expectation value of the evolution operator has the form obtained in theorems 1 and 2. Under assumptions

$$A = \lambda^2 A_2 = i\lambda^2 \int \frac{|v(k)|^2}{\omega(k)} d^d k < \infty, \quad B = \lambda^2 B_2 = -\lambda^2 \int \frac{|v(k)|^2}{\omega^2(k)} d^d k < \infty, \quad (3.28)$$

one has the following

Proposition 1. The vacuum expectation value of the evolution operator $U(t) = e^{itH_0} e^{-itH}$ is

$$\langle U(t) \rangle = \exp \left[At + B + \lambda^2 \int dk \frac{|v(k)|^2}{\omega^2(k)} e^{-i\omega(k)t} \right] \quad (3.29)$$

Proof. It follows from the explicit solution

$$H = W^*(H_0 + E_0)W, \quad E_0 = -\int \frac{|v(k)|^2}{\omega(k)} dk. \quad (3.30)$$

$$W = \exp \lambda \int \frac{(a^*(k)v(k) - a(k)\bar{v}(k))}{\omega(k)} dk = \exp \lambda \int \frac{a^*(k)v(k)}{\omega(k)} dk \exp -\lambda \int \frac{a(k)\bar{v}(k)}{\omega(k)} dk e^{B/2}$$

From Proposition 1 we obtain

Proposition 2. The asymptotic behaviour of the expectation value (3.29) for $t \rightarrow \infty$ has the form

$$\langle U(t) \rangle = \exp \left[At + B + \lambda^2 \left(\frac{1}{t} \right)^{\frac{d}{2}} (2i\pi)^{\frac{d}{2}} \frac{|v(k_0)|^2}{\omega^2(k_0)} e^{-i\omega(k_0)t} + \dots \right]. \quad (3.31)$$

where k_0 is a critical point, $\nabla\omega(k_0) = 0$ and we assume there is only one nondegenerate critical point.

Proof. It follows immediately from (3.29) by using the stationary phase method.

Remark 1. After the rescaling $t \rightarrow t/\lambda^2$ we get from (3.31) corrections to the stochastic limit

$$\langle U(t/\lambda^2) \rangle = \exp \left[A_2 t + B + \lambda^2 \left(\frac{\lambda^2}{t} \right)^{\frac{d}{2}} (2i\pi)^{\frac{d}{2}} \frac{|v(k_0)|^2}{\omega^2(k_0)} \cdot e^{-i\omega(k_0)t/\lambda^2} + \dots \right]. \quad (3.32)$$

Remark 2. One can take for example

$$v(k) = \omega(k) f(k),$$

$$\omega(k) = \frac{k^2}{2} - \omega_0, \quad \omega_0 > 0$$

where $f(k)$ is a test function. Then the assumptions (3.28) are satisfied,

$$B = -\lambda^2 \int |f(k)|^2 d^d k, \quad A = i\lambda^2 \int \left(\frac{k^2}{2} - \omega_0 \right) |f(k)|^2 d^d k,$$

the critical point $k_0 = 0$ and (3.32) has the form

$$\langle U(t) \rangle = \exp \left[At + B + \lambda^2 \left(\frac{1}{t} \right)^{\frac{d}{2}} (2i\pi)^{\frac{d}{2}} |f(0)|^2 e^{i\omega_0 t} + \dots \right]$$

Remark 3. If $\omega(k) = k^2 - \omega_0$ and $v(k)$ is an arbitrary test function then one has the decay. We can not use in this case the diagonalization of the Hamiltonian (3.27) but the formula (3.1) still is true. We have

$$\langle U(t) \rangle = \exp[-\lambda^2 \int_0^t dt_1 \int_0^{t_1} dt_2 \int d^d k |v(k)|^2 e^{i(t_2 - t_1)\omega(k)}] = \exp[\lambda^2 t A_2(t) + \lambda^2 B_2(t)]$$

where

$$A_2(t) = - \int_0^t d\sigma F(\sigma), \quad B_2(t) = \int_0^t d\sigma \sigma F(\sigma)$$

and

$$F(\sigma) = \int d^d k |v(k)|^2 e^{-i\sigma\omega(k)}$$

By using this representation and the expansion

$$F(\sigma) = \left(\frac{2i\pi}{\sigma} \right)^{\frac{d}{2}} |v(0)|^2 e^{i\sigma\omega(0)} \left(1 + o\left(\frac{1}{\sigma}\right) \right)$$

we compute corrections to the stochastic limit. Notice also that the stochastic limit

$$\lim_{\lambda \rightarrow 0} \lambda^2 \int_0^t dt_1 \int_0^{t_1} dt_2 F(t_1 - t_2) = \lim_{\lambda \rightarrow 0} \int_0^t d\tau \int_{-\tau/\lambda^2}^0 d\sigma F(-\sigma) = t \int_0^\infty d\sigma F(\sigma)$$

exists in any dimension $d = 1, 2, \dots$

3.4 Higher orders

To formulate the theorem about the asymptotic behaviour of the vacuum expectation value of the evolution operator it is convenient to use the Friedrichs Γ - operation

$$\Gamma(V) = \lim_{\epsilon \rightarrow +0} (-i) \int_0^\infty e^{-\epsilon t} e^{itH_0} V e^{-itH_0} dt \quad (3.33)$$

acting to the monomials as

$$\Gamma(V_{I,J}) = \int (\Gamma v)(p_1, i_1 \dots p_I, i_I | q_1, j_1 \dots q_J, j_J) \prod_{l=1}^I a_{i_l}^*(p_l) dp_l \prod_{r=1}^J a_{j_r}(q_r) dq_r \quad (3.34)$$

where

$$(\Gamma v)(p_1, i_1 \dots p_I, i_I | q_1, j_1 \dots q_J, j_J) = \frac{v(p_1, i_1 \dots p_I, i_I | q_1, j_1 \dots q_J, j_J)}{\sum_{l=1}^I \omega_l(p_l) - \sum_{r=1}^J \omega_r(p_r) + i0}.$$

It has the property

$$[H_0, \Gamma_\pm(V)] = V. \quad (3.35)$$

The following theorem gives a very effective exact representation for the vacuum expectation value of the evolution operator.

Theorem 3. Vacuum expectation value of the evolution operator for the Hamiltonian (2.3) has the following representation

$$\langle U(t) \rangle = e^{At+B+C(t)} \quad (3.36)$$

where

$$A = i \sum_{n=2}^\infty \lambda^n \langle 0 | \underbrace{(V \Gamma(V \dots \Gamma(V) \dots))}_n \rangle_c | 0 \rangle, \quad (3.37)$$

$$B = \sum_{n=2}^\infty \lambda^n \sum_{l=2}^\infty (-1)^{l-1} \sum_{k_1 + \dots + k_l = n} \langle 0 | (W_{k_1} \dots W_{k_l})_c | 0 \rangle, \quad (3.38)$$

and

$$W_n = \underbrace{\Gamma(V \dots \Gamma(V) \dots)}_n \quad (3.39)$$

The function $C(t)$ is given by the sum of terms of the form

$$\int dp \dots e^{it \sum_{l=1}^{k_1} E_k} \underbrace{\Gamma(v_{i1} \circ \dots \Gamma(v_{i2} \circ) \dots)}_{k_1} \underbrace{\Gamma(v_{i3} \circ \dots \Gamma(v_{i4} \circ) \dots)}_{k_2} \dots \underbrace{\Gamma(v_{i5} \circ \dots \Gamma(v_{i6} \circ) \dots)}_{k_{l+1}}(p, \dots). \quad (3.40)$$

To prove this Theorem let us consider one of the terms representing the n -th order of the perturbation theory with adiabatic parameter $\epsilon > 0$

$$\int dp \dots \mathcal{E}_0^{(n)}(p, \dots) = \quad (3.41)$$

$$(-i\lambda)^n \int dp \dots \int_0^t dt_1 \int_0^{t_1} dt_2 \dots \int_0^{n-1} dt_n e^{\sum_k (iE_k t_k - \epsilon t_k)} (v_{i1} \circ \dots \circ v_{in})(p \dots),$$

Integrating over t_n one gets two terms. One contains $e^{iE_n t_{n-1}}$ and the other contains 1. Together with the factor coming from $V(t_{n-1})$ the exponent $e^{iE_n t_{n-1}}$ yields to $e^{i\mathcal{E}_{n-1} t_{n-1}}$, where $\mathcal{E}_{n-1} = E_n + E_{n-1}$ is the total energy of the lines crossed by $n-1$ -cut (see Fig.2).




Figure 2: Diagram with energy-cuts

Therefore, the contribution coming from the upper bounds of integration over $t_n, \dots, t_2$ is equal to $e^{i\mathcal{E}_2 t_1}$, where $\mathcal{E}_2 = \sum_{k=2}^n E_k$. It is evident that $\mathcal{E}_2 = -E_1$ and therefore the integrand does not depend on t_1 and intergration of this terms over t_1 produces

$$t \int dp \dots \underbrace{(v \circ \Gamma(v \circ \dots \Gamma(v \circ) \dots))}_n(p \dots) \quad (3.42)$$

All these terms correspond to

$$t \langle 0 | \underbrace{(V \Gamma(V \dots \Gamma(V) \dots))}_n \rangle_c | 0 \rangle. \quad (3.43)$$

Or, in other words, we have proved that contributions from the upper bounds produce the linear in t terms. To prove that the term (3.43) gives the leading term in asymptotic expansion we have to prove that the rest terms are of order 1 when $t \rightarrow \infty$. Or in other words we have to

prove that the terms which contain at least one contribution from the down bound are of order 1.

To make more clear the proof we present in Appendix A an explicit calculation of $\mathcal{E}_0$ up to the fourth order of the perturbation theory.

Let us first consider contribution from only one the lower bound. For example, the integration over t_1 of the term produced by the low bound of integration over t_{k+1} and all other upper bounds gives:

$$\begin{aligned} & \int dp \dots \int_0^{t_1} dt_1 e^{i \sum_{l=1}^k E_l t_1} \underbrace{(v \circ \Gamma(v \circ \dots \Gamma(v \circ) \dots))}_k \underbrace{\Gamma(v \circ \dots \Gamma(v) \dots))}_{n-k} (p \dots) = \\ & i \int dp \dots [e^{it \sum_{l=1}^k E_l} - 1] \underbrace{(\Gamma(v \circ \Gamma(v \circ \dots \Gamma(v \circ) \dots))}_k \underbrace{\Gamma(v \circ \dots \Gamma(v) \dots))}_{n-k} (p \dots). \end{aligned} \quad (3.44)$$

We see that (3.44) gives a contribution to terms in (3.2) that do not depend on t . All these contributions from the down bound of integration over t_k and t_1 and all other upper bounds sum up to the following expression

$$- \langle 0 | \underbrace{(\Gamma V \Gamma(V \dots \Gamma(V) \dots))}_k \underbrace{\Gamma(V \dots \Gamma(V) \dots))}_{n-k} | 0 \rangle. \quad (3.45)$$

Let us note the obvious difference between

$$\underbrace{\Gamma(V \dots \Gamma(V) \dots)}_k \underbrace{\Gamma(V \dots \Gamma(V) \dots)}_l \quad \text{and} \quad \underbrace{\Gamma(V \dots \Gamma(V) \dots)}_{k+l}. \quad (3.46)$$

This expression can be presented on graphs, see Fig.3. On the graph the directions of applications of Γ operators are indicated by brackets. The term in (3.44) with $e^{it \sum_{l=1}^k E_l}$ produces the decreasing contribution as $t \rightarrow \infty$ since we have $\sum_{l=1}^k E_l > 0$.




Figure 3: Diagram corresponding to $\Gamma(V\Gamma(V\Gamma(V)))\Gamma(V\Gamma(V\Gamma(V)))$

For the n -th order of perturbation theory the terms corresponding to contributions of l lower bounds have the following structure

$$\int dp \dots [e^{it \sum_{k=1}^{k_1} E_k} - 1] \underbrace{\Gamma(v \circ \dots \Gamma(v \circ) \dots)}_{k_1} \underbrace{\Gamma(v \circ \dots \Gamma(v \circ) \dots)}_{k_2} \dots \underbrace{\Gamma(v \circ \dots \Gamma(v) \dots)}_{k_{l+1}}(p, \dots) \quad (3.47)$$

$\sum_{i=1}^{l+1} k_i = n$. All these terms either contain oscillator factors or give the constant factors. The constant factors can be sum up to expressions

$$(-)^l \langle 0 | \underbrace{\Gamma(V \dots \Gamma(V) \dots)}_{k_1} \underbrace{\Gamma(V \dots \Gamma(V) \dots)}_{k_2} \dots \underbrace{\Gamma(V \dots \Gamma(V) \dots)}_{k_{l+1}} \rangle_c | 0 \rangle. \quad (3.48)$$

Therefore we have shown that

$$\mathcal{E}_0 = tA + B + C(t), \quad (3.49)$$

where A is given in perturbation theory by (3.37) and B is given by

$$B = \sum_n \lambda^n \sum_l \sum_{k_1 + \dots + k_l = n} (-1)^{l-1} \langle 0 | \underbrace{\Gamma(V \dots \Gamma(V) \dots)}_{k_1} \underbrace{\Gamma(V \dots \Gamma(V) \dots)}_{k_2} \dots \underbrace{\Gamma(V \dots \Gamma(V) \dots)}_{k_l} \rangle_c | 0 \rangle. \quad (3.50)$$

The function $C(t)$ is the sum of terms of the form

$$\int dp \dots e^{it \sum_{l=1}^{k_1} E_k} \underbrace{\Gamma(v \circ \dots \Gamma(v \circ) \dots)}_{k_1} \underbrace{\Gamma(v \circ \dots \Gamma(v \circ) \dots)}_{k_2} \dots \underbrace{\Gamma(v \circ \dots \Gamma(v) \dots)}_{k_{l+1}}(p, \dots). \quad (3.51)$$

These terms vanish as $t \rightarrow \infty$ because $\sum_{l=1}^{k_1} E_k \neq 0$.

3.5 Wave operators and the main formula

In this section we show how the spectral theory and renormalized wave operators can be used for the derivation of the main formula. In particular explicit expressions for the parameters A, B and $C(t)$ will be obtained.

We consider the Hamiltonian (2.3) for one type of particles with $\omega(p) = \sqrt{p^2 + m^2}$, $m > 0$ in the space R^d , $d > 2$. We will work with the formal perturbation series for the evolution operator. In fact if the interaction (2.3) includes only fermionic operators or it is linear in bosonic operators then one can prove that the series are absolutely convergent. The following operator plays the crucial role in the scattering theory

$$T =: \exp \sum_{n=1}^{\infty} (-\lambda)^n \Gamma(V \dots \Gamma(V) \dots)_L : \quad (3.52)$$

Here $()_L$ means that only connected non-vacuum diagrams are included. The operator T is equal in fact to the non-vacuum part of the conjugate wave operator :

$$T = \lim_{\epsilon \rightarrow 0} \lim_{t \rightarrow \infty} U_{\epsilon}^*(t) / \langle U_{\epsilon}^*(t) \rangle$$

One has the following relations [5]:

$$HT = T(H_0 + E_0), \quad (3.53)$$

$$E_0 = \sum_{n=1}^{\infty} (-\lambda)^{n+1} < V\Gamma(V\ldots\Gamma(V)\ldots)_c >,$$

$$T^*T = TT^* = Z^{-1},$$

$$Z^{-1} = ||T\Phi_0||^2$$

We will use these relations to derive the main formula for matrix elements of the evolution operator and in particular to compute corrections to the stochastic limit.

From (3.53) it follows

$$H = T(H_0 + E_0)T^*e^B$$

where

$$B = \ln Z \quad (3.54)$$

Therefore one has

$$U(t) = e^{itH_0}e^{-itH} = e^{itH_0}Te^{-it(H_0+E_0)}T^*e^B = e^{At+B}e^{itH_0}Te^{-itH_0}T^* \quad (3.55)$$

where

$$A = -iE_0 \quad (3.56)$$

By taking the expectation value of the equality (3.55) we obtain

$$< \psi, U(t)\psi > = e^{At+B+C(t)} \quad (3.57)$$

where

$$e^{C(t)} = < \psi, e^{itH_0}Te^{-itH_0}T^*\psi >$$

If ψ is the vacuum vector then one can prove that $C(t) \rightarrow 0$ as $t \rightarrow \infty$ and we obtain the main formula (3.36). If ψ is a non-vacuum vector then the asymptotic behaviour of $C(t)$ is more complicated. We have proved the following theorem.

Theorem 4. If the Hamiltonian satisfies the indicated above assumptions then there exists the following representation for the vacuum expectation value of the evolution operator

$$< \Phi_0, U(t)\Phi_0 > = e^{At+B+C(t)}$$

where constants A and B are given by (3.50) and (3.38) and $C(t)$ is defined as

$$e^{C(t)} = < \Phi_0, T(t)T^*\Phi_0 >, \quad T(t) = e^{itH_0}Te^{-itH_0} \quad (3.58)$$

here the weak limit of $T(t)$ as $t \rightarrow \infty$ is equal to 1 and $\lim_{t \rightarrow \infty} C(t) = 0$.

This theorem is closely related with the theorem 3 and it shows the physical meaning of constants A, B and the function $C(t)$.

4 One particle matrix element of the evolution operator for translation invariant Hamiltonian

In this section we study the asymptotic behaviour of one particle matrix elements of evolution operator

$$\langle p|U(t, \lambda)|p' \rangle = \delta(p - p')U_{1,1}(t, p, \lambda), \quad (4.1)$$

for translation invariant Hamiltonian (2.4) without vacuum polarization, i.e. when $V_{I0} = V_{0J} = 0$, and we assume also $V_{1,1} = 0$. We will prove the following

Theorem 5. For $t\lambda^2 = \text{fixed} = \tau$ and small λ^2 one has the following representation

$$U_{1,1}(t, p, \lambda) = \exp\{i\tau A_2(p)\}(1 + i\lambda^2\tau A_4(p) + \lambda^2 B_2(p) + o(\lambda^2)) \quad (4.2)$$

where

$$A_2(p) = (V\Gamma(V))_{1,1}(p) \quad (4.3)$$

$$B_2(p) = (V\Gamma^2(V))_{1,1}(p) \quad (4.4)$$

$$A_4(p) = A_4^{1PI}(p) + A_4^{1PR}(p),$$

$$A_4^{1PI}(p) = (V\Gamma(V\Gamma(V\Gamma(V))))_{1,1}^{1PI}(p) \quad (4.5)$$

$$A_4^{1PR}(p) = (V\Gamma(V)(V\Gamma^2(V)))_{1,1}^{1PR}(p) \quad (4.6)$$

Here subscript $_{1,1}$ means that we left only connected diagrams in the corresponding expression with one creations and one annihilation operator and we take only the coefficient in front of $\delta(p - p')$,

$$\langle p|\mathcal{O}|p' \rangle = \mathcal{O}_{1,1}(p)\delta(p - p') \quad (4.7)$$

Remark 1. We will give the proof of Theorem 5 for the Hamiltonian of the form

$$H = \int \omega(p)a^+(p)a(p)dp + \lambda \int (v(p, q)a^+(q)a^+(p - q)a(p) + c.c.)dpdq. \quad (4.8)$$

Generalization to the Hamiltonians in the general form (2.4) is obvious and the suitable comments will be given below.

Remark 2. In fact the following representation is true

$$U_{1,1}(t, p, \lambda) = \exp\{itA(p, \lambda) + B(p, \lambda)\}(1 + C(t, p, \lambda)) \quad (4.9)$$

where $A(p, \lambda)$, $B(p, \lambda)$ and $C(t, p, \lambda)$ are formal series in λ

$$A(p, \lambda) = \sum_{n=1}^{\infty} \lambda^n A_n(p), \quad B(p, \lambda) = \sum_{n=1}^{\infty} \lambda^n B_n(p), \quad C(t, p, \lambda) = \sum_{n=1}^{\infty} \lambda^n C_n(t, p) \quad (4.10)$$

and functions $C_n(t, p)$ vanish as $t \rightarrow \infty$. In the case of $\langle p|V|p' \rangle = 0$, $A_1 = 0$, $B_1 = 0$ and low order terms in these series are given by (4.3), (4.4) and

$$C_2(t, p) = (e^{itH_0} V e^{-itH_0} \Gamma^2(V))_{1,1}(p). \quad (4.11)$$

We will give the proof of (4.2) in the next section using the methods of scattering theory. This proof assumes that one has not decay.

Remark 3. Representation (4.2) means a special dependence of $U_{1,1}(t, \lambda, p)$ on t . Namely it says that (2n)-order on λ of $U_{1,1}(t, \lambda, p)$

$$U_{1,1}(t, \lambda, p) = \sum_n \lambda^{2n} U_{1,1}^{(2n)}(t, p),$$

can be represent as

$$U_{1,1}^{(2n)}(t, p) = \sum_{k=0}^n t^k U_{1,1}^{(2n,k)}(p) + r_n(t, p) \quad (4.12)$$

with $r_n(t) \rightarrow 0$ for $t \rightarrow \infty$.

Remark 4. Performing the stochastic limit we get

$$U_{1,1}(\tau/\lambda^2, p) \rightarrow \exp i\tau A_2(p) \text{ as } \lambda \rightarrow 0, \quad (4.13)$$

i.e. the stochastic limit of one particle matrix elements of evolution operator, $U(\tau/\lambda^2)$, for translation invariant Hamiltonian without vacuum polarization is given by the simple second order diagram (Fig.9). This result takes place for arbitrary interaction (2.4) and the answer can be representing in the form

$$U_{1,1}(\tau/\lambda^2, p) \rightarrow \exp\{i\tau(VT(V))_{1,1}^{(c)}(p)\} \text{ as } \lambda \rightarrow 0. \quad (4.14)$$

Here the symbol $(\dots)_{1,1}^{(c)}$ means that we left only connected diagrams in $VT(V)$ with one creations and one annihilation operator corresponding to the same type of particles.




Figure 4: The second order mass renormalization diagram

Remark 5. There are two sources of corrections to the stochastic limit. One is the terms in the second order approximation of $U_{1,1}^{(2)}(t)$ which does not depend on t , i.e. it corresponds to diagram Fig.9. There are also corrections to the leading terms of the stochastic limit which

come from 4-th order terms with linear dependence on t . These corrections come from the 4-th order 1PI diagrams such as shown on Fig.5. (see for details Appendix where we show that contributions of these diagrams include the terms proportional to t and after rescaling produce a factor $\tau\lambda^2$).

$$U_{1,1}(\tau/\lambda^2, p) \rightarrow \exp\{i\tau A_2(p)\}[1 + \lambda^2(i\tau A_4(p)it + B_2(p))] \text{ as } \lambda \rightarrow 0. \quad (4.15)$$




Figure 5: The 4-order 1PI mass renormalization diagrams

Remark 6. Let us note that $A_2(p)$ in (4.3) is positive in the case when we assume that there is no decay in the model. For Hamiltonian (4.8) the explicit expression for $A_2(p)$ is

$$A_2(p) = \int dq (\bar{v} \circ \Gamma v)(p, q), \quad (4.16)$$

where

$$(\bar{v} \circ \Gamma v)(p, q) = \frac{|v(p, q)|^2}{\omega(p - q) + \omega(q) - \omega(p)} \quad (4.17)$$

We see that for the typical example $\omega(q) = \sqrt{q^2 + m^2}$ the denominator in (4.17) does not vanish. In the case of different particles, say, 3 different particles (Lee model)

$$H = \int \omega_a(p) a^+(p) a(p) dp + \int \omega_b(p) b^+(p) b(p) dp + \int \omega_c(p) c^+(p) c(p) dp + \lambda \int dp dq (v(p, q) (a^+(q) b^+(p - q) c(p) + c.c.)) \quad (4.18)$$

we have the ϵ -prescription for the denominator

$$A_2(p) = \int dq \frac{|v(p, q)|}{\omega_a(p - q) + \omega_b(q) - \omega_c(p) + i\epsilon} \quad (4.19)$$

and if one has a decay then $A_2(p)$ gets the imaginary part.

Proof. Now let us prove the Theorem 5. Let us first make calculations up to the second order ¹. There is one the second-order diagram (Fig.4) and it gives :

$$\mathcal{U}_2(t, p, q) = (-i\lambda)^2 \int_0^t dt_1 \int_0^{t_1} dt_2 |v(p, q)|^2 e^{iE_2(-t_1+t_2)} =$$

¹ An explicit calculation of $U_{1,1}(t, p, \lambda)$ up to 4-th order is presented in Appendix B




Figure 6: Reducible mass renormalization diagram with two irreducible mass insertions

$$it\lambda^2(\bar{v} \circ \Gamma v)(p, q) - \lambda^2(\bar{v} \circ \Gamma^2 v)(p, q)(1 - e^{iE_1 t}) \quad (4.20)$$

where $E_1 = -E_2$, $E_2 = \omega(k) + \omega(p - k) - \omega(p)$. Here and below for simplicity of writing only the kernel $\mathcal{U}_{1,1}(T, p, q_1, \dots)$ of $U_{1,1}(t, p)$ is written,

$$U_{1,1}^{(2n)}(t, p) = \int dq_1 \dots dq_r \mathcal{U}_{2n}(t, p, q_1, \dots q_r). \quad (4.21)$$

In the case of a general Hamiltonian instead of (4.20) one has

$$U_{1,1}^{(2)}(t, p) = it\lambda^2(V\Gamma(V))_{1,1}(p) - \lambda^2(V\Gamma^2(V))_{1,1}(p) + \lambda^2(e^{iH_0 t} V e^{-iH_0 t} \Gamma^2(V))_{1,1}(p). \quad (4.22)$$

We will give the proof by induction. We already has proved the claim of the theorem at the second order of perturbation theory. Let us suppose that the 1PR diagrams including n irreducible second order mass insertions (see Fig.7) have the following behaviour




Figure 7: 1PR mass renormalization diagram with n irreducible 2-order insertions

$$U_{1,1}^{1P(n)R}(t)(p) = \frac{(it)^n}{n!} A_2^n(p) + \frac{n(it)^{n-1}}{(n-1)!} B_2(p) A_2^{n-1}(p) + \dots \quad (4.23)$$

Here down index means the order in λ and index n in upper script means the number of irreducible insertions and dots means low order on T .

We also assume that the 1PR diagrams which include $(n-2)$ irreducible second order mass insertions and one irreducible 4-th order mass insertion (see Fig.8) have the following behaviour

$$U_{2n}^{1P(n-1)R}(t)(p) = \frac{(it)^{n-1}}{(n-2)!} A_4^{1PI}(p) A_2^{n-2}(p) + \dots \quad (4.24)$$




Figure 8: 1PR mass renormalization diagram with $(n-1)$ irreducible 2-order insertions and one irreducible 4-order insertion

1RR diagrams with $(n+1)$ irreducible second order mass insertions (see Fig.7) have the following representation

$$U_{2(n+1)}^{1P(n+1)R}(t, p) = (-i)^2 \int_0^t dt_1 \int_0^{t_1} dt_2 \int dq |v(p, q)|^2 e^{iE_2(-t_1+t_2)} U_{2n}^{1P(n)R}(t_2, p), \quad (4.25)$$

Using the assumption (4.23) and

$$\int_0^t dt_1 \int_0^{t_1} dt_2 e^{iE(-t_1+t_2)} t_2^n = \frac{t^{n+1}}{(n+1)(iE)} - \frac{t^n}{(iE)^2} + \frac{n(n-1)}{n+1} \frac{t^{n-1}}{(iE)^3} + \dots \quad (4.26)$$

we see that the RHS of (4.25) has the following behaviour

$$\begin{aligned} & (-i)^2 \int_0^t dt_1 \int_0^{t_1} dt_2 \int dq |v(p, q)|^2 e^{iE_2(-t_1+t_2)} \left[\frac{(it_1)^n}{n!} A_2^n(p) + \frac{n(it_1)^{n-1}}{(n-1)!} B_2(p) A_2^{n-1}(p) + \dots \right] = \\ & (i)^2 \int dq \left\{ \frac{(it)^{n+1}}{i(n+1)!} \frac{|v(p, q)|^2}{iE_2} A_2^n(p) - \frac{(it)^n}{n!} \frac{|v(p, q)|^2}{(iE_2)^2} A_2^n(p) + n \frac{(it)^n}{in!} \frac{|v(p, q)|^2}{iE_2} B_2(p, q) A_2^{n-1}(p) + \dots \right\}, \end{aligned}$$

i.e. the contribution of 1RR diagrams with $(n+1)$ irreducible second order parts (see Fig.7) has the following asymptotic

$$U_{2(n+1)}^{1P(n+1)R}(t, p) = \frac{(it)^{n+1}}{(n+1)!} A_2^{n+1}(p) + (n+1) \frac{(it)^n}{n!} B_2(p) A_2^n(p) + \dots \quad (4.27)$$

Here

$$A_2(p) = \int dq \frac{|v(p, q)|^2}{E_2}, \quad B_2(p) = - \int dq \frac{|v(p, q)|^2}{E_2^2}, \quad (4.28)$$

or in the more general case

$$A_2(p) = (VT(V))_{1,1}^{1PI}(p), \quad B_2(p) = -(VT^2(V))_{1,1}^{1PI}(p) \quad (4.29)$$

Therefore, (4.27) is in accordance with (4.23), i.e. (4.23) is proved by induction.

1PR diagrams included $(n-1)$ irreducible second order mass insertions and one irreducible 4-th order mass insertion (see Fig.8) have the following representation

$$U_{2(n+1)}^{1P(n)R}(t, p) = (-i)^2 \int_0^t dt_1 \int_0^{t_1} dt_2 \int dq |v(p, q)|^2 e^{iE_2(-t_1+t_2)} U_{2n}^{1P(n-1)R}(t_2, p) + \quad (4.30)$$

$$(-i)^4 \int_0^t dt_1 \int_0^{t_1} dt_2 \int_0^{t_2} dt_3 \int_0^{t_3} dt_4 \int dq dq_1 (\bar{v} \circ \bar{v} \circ v \circ v)_{1,1}^{(1PI)}(p, q, q_1) e^{i \sum_{i=1}^4 E_i t_i} U_{2(n-1)}^{1P(n-1)R}(t_4, p)$$

Using

$$\int_0^t dt_1 \int_0^{t_1} dt_2 \int_0^{t_2} dt_3 \int_0^{t_3} dt_4 t_4^n e^{i \sum_{i=1}^4 E_i t_i} = \frac{1}{i^3} \frac{t^{n+1}}{n+1} \frac{1}{(E_4)(E_4 + E_3)(E_4 + E_3 + E_2)}, \quad (4.31)$$

(4.26) and the assumptions (4.23) and (4.24) we see that the RHS of (4.30) has the following behaviour

$$(i)^2 \int_0^t dt_1 \int_0^{t_1} dt_2 \int dq |v(p, q)|^2 e^{i E_2 (-t_1 + t_2)} \left[\frac{(it_2)^{n-1}}{(n-2)!} A_4^{1PI}(p) A_2^{n-2}(p) + \dots \right] +$$

$$(i)^4 \int_0^t dt_1 \int_0^{t_1} dt_2 \int_0^{t_2} dt_3 \int_0^{t_3} dt_4 dq dq_1 (\bar{v} \circ \bar{v} \circ v \circ v)_{1,1}^{(1PI)}(p, q, q_1) e^{i \sum_{i=1}^4 E_i t_i} \left[\frac{(it_4)^{n-1}}{(n-1)!} A_2^{n-1}(p) + \dots \right] =$$

$$= \frac{(it)^n}{(n-2)!n} \int dq (\bar{v} \circ \Gamma(v))_{1,1}^{(1PI)}(p, q) A_4^{1PI} A_2^{n-2} + i \frac{(it)^n}{n!} \int dq dq_1 (\bar{v} \circ \Gamma(\bar{v} \circ \Gamma(v \Gamma(v))))_{1,1}^{(1PI)}(p, q, q_1) A_2^{n-1},$$

Here

$$A_4^{1PI}(p) = \int dq dq_1 (\bar{v} \circ \Gamma(\bar{v} \circ \Gamma(v \Gamma(v))))_{1,1}^{(1PI)}(p, q, q_1) \quad (4.32)$$

or more generally,

$$A_4^{1PI}(p) = (V \Gamma(V \Gamma(V \Gamma(V))))_{1,1}^{(1PI)}(p) \quad (4.33)$$

Therefore,

$$U_{2(n+1)}^{1P(n-1)R}(t, p) = \frac{(it)^n}{(n-1)!} t A_4(p) A_2^{n-1}(p),$$

that is in accordance with assumption (4.24).

It is evident that the behaviour (4.23) and (4.24) implies (4.2) with A_2 and B_2 as in the formulation of Theorem 5.

4.1 Wave operators and the main formula

In this section we show how the spectral theory and renormalized wave operators can be used for the derivation of the main formula. In particular, explicit recursive relations for the parameters A_n , B_n and $C_n(t)$ will be obtained. Note that for this derivation we have to assume that the is no decay.

The intertwining operator T is defined as a solution of the following equation

$$HT = T(H_0 + M), \quad (4.34)$$

Here M has the form

$$M = \int m(p) a^*(p) a(p) dp \quad (4.35)$$

The operator T plays the crucial role in the scattering theory. Its singular part defines the renormalized wave operators. The renormalized wave operators also give a solution of intertwining condition. Taking T in the following form

$$T =: \exp W :, \quad W = \Gamma(Q) \quad (4.36)$$

one gets [6] equations to define Q and M .

$$Q + V \perp T - (V \perp T)_{1,1} - W - \circ - M = 0 \quad (4.37)$$

$$M = (V \perp T)_{1,1} \quad (4.38)$$

Here the symbol $\perp$ means that for connected A in the operators $A \perp B$ all connected parts of B are paired with A . If B is connected then $A \perp B = A - \circ - B = (AB)_c$. For a special form of the interaction, when $V_{0I} = V_{I0} = 0$ we can write $M = (V \perp T)_{1,1} = (V\Gamma(Q))_{1,1}$.

Expanding M and Q in the power series in λ

$$M = \sum_{n=1} \lambda^{2n} M_{2n}, \quad Q = \sum_{n=1} \lambda^{2n+1} Q_{2n+1}, \quad (4.39)$$

we get recursive relations to define M_{2n} and Q_{2n+1} . Let us compute explicitly the first terms solving these equations. We obtain

$$Q_1 = -V \quad (4.40)$$

$$M_2 = -(V\Gamma V)_{1,1} \quad (4.41)$$

$$Q_2 = (V\Gamma V)_c - (V\Gamma V)_{1,1} \quad (4.42)$$

$$Q_3 = -(V\Gamma_r(V\Gamma(V)))_c - \frac{1}{2}V \perp : \Gamma(V)^2 : + \Gamma V - \circ - M_2 \quad (4.43)$$

$$M_4 = -(V\Gamma(V\Gamma_r(V\Gamma(V))))_{1,1} + (V\Gamma^2(V)M_2)_{1,1} \quad (4.44)$$

or

$$M_4 = -(V\Gamma(V\Gamma_r(V\Gamma(V))))_{1,1} - (V\Gamma^2(V)(V\Gamma(V))_{1,1})_{1,1} \quad (4.45)$$

Here we use the notation

$$\Gamma_r(Q) = \Gamma(Q - Q_{1,1}) \quad (4.46)$$

One can construct in perturbation theory the operator Z such that

$$TT^*Z = 1 \quad (4.47)$$

In the second order in λ

$$\langle p|Z|p' \rangle = (1 + \lambda^2 B_2(p))\delta(p - p') \quad (4.48)$$

Therefore one has

$$U(t) = e^{itH_0} e^{-itH} = e^{-itM} e^{it(H_0+M)} T e^{-it(H_0+M)} T^* Z \quad (4.49)$$

By taking the one particle expectation value of the equality (4.49)

$$\langle p|U(t)|p' \rangle = e^{-iM(p,\lambda)t} Z(p, \lambda) (1 + C(p, t, \lambda)) \quad (4.50)$$

where

$$(1 + C(t, p, \lambda))\delta(p - p') = \langle p | e^{it(H_0+M)} T e^{-it(H_0+M)} T^* | p' \rangle \quad (4.51)$$

and M and T should be computed from the recursive relations.

We have proved the following

Theorem 6. For translation invariant Hamiltonians without vacuum polarization the one particle matrix elements of evolution operator are given by the formula (4.50) where functions $M(p, \lambda)$ and $Z(p, \lambda)$ are solutions of equations (4.36)-(4.38) and (4.47). Function $C(t, p, \lambda)$ is defined by (4.51) and it vanishes as $t \rightarrow \infty$.

5 Conclusion

We have obtained in this paper the explicit representations for the vacuum and one-particle matrix elements of the evolution operator. By using these representations we have computed the corrections to the known results for the large time exponential behaviour of these matrix elements. This opens the way for further investigations of the large time behaviour in quantum theory. In particular the problems of quantum decoherence and decay require the further study by using these methods.

Acknowledgments. This work is supported in part by INTAS grant 96-0698, I.Ya.A. is supported also by RFFI-99-01-00166 and I.V.V. by RFFI-99-01-00105 and by grant for the leading scientific schools.

APPENDIX

A 4-th order for hamiltonian with vacuum polarization


For an illustration of the form (3.47) of vacuum expectation value of evolution operator we present on Fig.9 all contributions to the 4-th order of perturbation theory for vacuum expectation value of evolution operator. For simplicity we consider the interaction in the form

$$V = \int (v(p_1, p_2 p_3)(a^*(p_1) + a(p_1))((a^*(p_2) + a(p_2))a^*(p_3) + a(p_3))dp_1 dp_2 dp_3 \quad (A.1)$$

On this example we see once again that only one term from all terms represented on Fig.9 has linear dependens on t . This term is represented by bracket-diagram on which all right bracked are on the right from the last vertex.


Diagrams

Corresponding order of application of Γ -operator,
exponential factors and denominators:




$$T < 0 | V \Gamma(V \Gamma(V \Gamma(V))) | 0 > =$$

$$T \int dp \dots \frac{(\bar{v} \circ \bar{v} \circ v \circ v)(p, \dots)}{(E_1 + E_2 + E_3 + E_4)(E_2 + E_3 + E_4)(E_3 + E_4)E_4}$$




$$- \int dp \dots [e^{i(E_1 + E_2 + E_3)T} - 1] \cdot \Gamma(v \circ \Gamma(v \circ) \Gamma(v \circ)) \Gamma(v)(p, \dots) =$$

$$- \int dp \dots [e^{i(E_1 + E_2 + E_3)T} - 1] \frac{(\bar{v} \circ \bar{v} \circ v \circ v)(p, \dots)}{(E_1 + E_2 + E_3 + i0)(E_2 + E_3 + i0)(E_3 + i0)E_4}$$




$$- \int dp \dots [e^{i(E_1 + E_2)T} - 1] \cdot \Gamma(\bar{v} \circ \Gamma(\bar{v} \circ)) \Gamma(v \circ \Gamma(v))(p, \dots)$$


$$- \int dp \dots [e^{i(E_1 + E_2)T} - 1] \cdot \frac{(\bar{v} \circ \bar{v} \circ v \circ v)(p, \dots)}{(E_1 + E_2 + i0)(E_2 + i0)(E_3 + E_4 + i0)E_4}$$




$$\begin{aligned}
& - \int dp \dots [e^{i(E_1)T} - 1] \Gamma(\bar{v} \circ) \Gamma(\bar{v} \circ \Gamma(\bar{v} \circ \Gamma(v))) (p, \dots) = \\
& - \int dp \dots [e^{i(E_1)T} - 1] \cdot \frac{(\bar{v} \circ \bar{v} \circ v \circ v)(p, \dots)}{(E_1 + i0)(E_2 + E_3 + E_4 + i0)(E_3 + E_4 + i0)E_4}
\end{aligned}$$




$$\begin{aligned}
& - \int dp \dots [e^{i(E_1 + E_2)T} - 1] \cdot \Gamma(\bar{v} \circ \Gamma(\bar{v} \circ)) \Gamma(v \circ) \Gamma(v) (p, \dots) = \\
& - \int dp \dots [e^{i(E_1 + E_2)T} - 1] \cdot \frac{(\bar{v} \circ \bar{v} \circ v \circ v)(p, \dots)}{(E_1 + E_2 + i0)(E_2 + i0)(E_3 + i0)E_4}
\end{aligned}$$



$$\begin{aligned}
& \int dp \dots [e^{iE_1 T} - 1] \cdot |\Gamma(\bar{v} \circ) \Gamma(\bar{v} \circ \Gamma(v \circ)) \Gamma(v) (p, \dots) = \\
& \int dp \dots [e^{iE_1 T} - 1] \cdot \frac{(\bar{v} \circ \bar{v} \circ v \circ v)(p, \dots)}{E_1(E_2 + E_3 + i0)(E_3 + i0)E_4}
\end{aligned}$$



$$\begin{aligned}
& \int dp \dots [e^{iE_1 T} - 1] \cdot \Gamma(\bar{v} \circ) \Gamma(\bar{v} \circ) \Gamma(v \circ \Gamma(v)) (p, \dots) \\
& \int dp \dots [e^{iE_1 T} - 1] \cdot \frac{(\bar{v} \circ \bar{v} \circ v \circ v)(p, \dots)}{E_1(E_2 + i0)(E_3 + E_4 + i0)E_4}
\end{aligned}$$



$$\begin{aligned}
& - \int dp \dots [e^{iE_1 T} - 1] \cdot \Gamma(\bar{v} \circ) \Gamma(\bar{v} \circ) \Gamma(v \circ) \Gamma(v)(p, \dots) \\
& - \int dp \dots [e^{iE_1 T} - 1] \cdot \frac{(\bar{v} \circ \bar{v} \circ v \circ v)(p, \dots)}{E_1(E_2 + i0)(E_3 + i0)E_4}
\end{aligned}$$

Figure 9: 4-th order diagrams of vacuum expectation value of evolution operator for interaction (A.1)

B 4-th order of one particle expectation value of evolution operator for translation invariant interaction

Here we prove the representation (4.2) - (4.6) for one particle matrix elements of evolution operator for translation invariant Hamiltonian (2.4) without vacuum polarization in the 4-th order of perturbation theory. We obtain

$$A_4(p) = A_4^{1PI}(p) + A_4^{1PR}(p),$$

$$A_4^{1PI}(p) = (V\Gamma(V\Gamma(V\Gamma(V))))_{1,1}^{1PI}(p), \quad (\text{B.1})$$

$$A_4^{1PR}(p) = (V\Gamma(V)(V\Gamma^2(V)))_{1,1}^{1PR}(p) \quad (\text{B.2})$$

There are two one-particle irreducible (1PI) diagrams (Fig. 5) and one one-particle reducible (1PI) diagram, (Fig. 6). The contribution of the 4-th order 1PI-diagram is

$$\begin{aligned}
\mathcal{U}_4^{1PR}(t, p, q, k) &= (-i\lambda)^4 \int_0^T dt_1 \int_0^{t_1} dt_2 \int_0^{t_2} dt_3 \int_0^{t_3} dt_4 |v(p, q)|^2 \cdot |v(p, k)|^2 e^{iE_1(t_1 - t_2)} e^{iE_4(t_4 - t_3)} = \\
& (-i\lambda)^2 \int_0^t dt_1 \int_0^{t_1} dt_2 |v(p, k)|^2 e^{iE_1(t_1 - t_2)} \mathcal{U}_2(t, p, q)
\end{aligned}$$

Substituting here (4.20) we have

$$\mathcal{U}_4^{1PR}(t)(p, q, k) = \frac{(iT)^2}{2} \mathcal{A}_2(p, q) \mathcal{A}_2(p, k) \quad (\text{B.3})$$

$$-iT [B_2(p, q) \mathcal{A}_2(p, k) + \mathcal{A}_4^{(1PR)}(p, q, k)] + \mathcal{B}_4^{1PR}(p, q, k) + \mathcal{C}_4^{1PR}(t, p, q, k)$$

where

$$\mathcal{A}_2(p, q) = (\bar{v} \circ \Gamma(v))(p, q) \quad (\text{B.4})$$

$$\mathcal{A}_4^{(1PR)}(p, q, k) = (\bar{v} \circ \Gamma(v))(p, q) (\bar{v} \circ \Gamma^2(v))(p, k) \quad (\text{B.5})$$

$$\begin{aligned} \mathcal{B}_4^{(1PR)}(p, q, k) = & [(\bar{v} \circ \Gamma^3(v))(p, k)(\bar{v} \circ \Gamma(v))(p, q) + (\bar{v} \circ \Gamma^2(v))(p, k)(\bar{v} \circ \Gamma^2(v))(p, q)] + \\ & (\bar{v} \circ \Gamma(\Gamma^2(v) \circ \bar{v}) \circ \Gamma(v))(p, q, k) + (\bar{v} \circ \Gamma(\Gamma(v) \circ \bar{v}) \circ \Gamma^2(v))(p, q, k) \end{aligned} \quad (\text{B.6})$$

$$\begin{aligned} C_4^{(1PR)}(t, p, q, k) = & -e^{iE_1 T} [(\bar{v} \circ \Gamma^3(v))(p, k)(\bar{v} \circ \Gamma(v))(p, q) + (\bar{v} \circ \Gamma^2(v))(p, k)(\bar{v} \circ \Gamma^2(v))(p, q)] \\ & e^{iE_1 t} (\bar{v} \circ \Gamma(\Gamma^2(v) \circ \bar{v}) \circ \Gamma(v))(p, q, k) + e^{iE_3 t} (\bar{v} \circ \Gamma(\Gamma(v) \circ \bar{v}) \circ \Gamma^2(v))(p, q, k), \end{aligned} \quad (\text{B.7})$$

here $E_1 = \omega(p) - \omega(q) - \omega(p - q)$ and $E_3 = \omega(p) - \omega(k) - \omega(p - k)$.

(B.3) can be rewritten in more compact form

$$U_{1,1}^{1PR}(t, p) = \frac{(it)^2}{2} A_2^2(p) - it [B_2(p)A_2(p) + A_4^{(1PR)}(p)] + B_4^{1PR}(p) + C_4^{1PR}(T, p) \quad (\text{B.8})$$

where

$$A_2(p) = (V\Gamma(V))_{1,1}(p), \quad A_4^{(1PR)}(p) = (V\Gamma(V))_{1,1}(p)(V\Gamma^2(V))_{1,1}(p) \quad (\text{B.9})$$

$$B_4^{(1PR)}(p) = (V\Gamma^3(V))_{1,1}(p)(V\Gamma(V))_{1,1}(p) + (V\Gamma^2(V))_{1,1}(p)(V\Gamma^2(V))_{1,1}(p) + \quad (\text{B.10})$$

$$(V\Gamma(\Gamma^2(V)V)\Gamma(V))_{1,1}^{1PR}(p) + (V\Gamma(\Gamma(V)V)\Gamma^2(V))_{1,1}^{1PR}(p)$$

$$C_4^{(1PR)}(t, p) = -(e^{iH_0 T} V\Gamma^3(V))_{1,1}(p)(V\Gamma(V))_{1,1}(p) - (e^{iH_0 t} V\Gamma^2(V))_{1,1}(p)(V\Gamma^2(V))_{1,1}(p) +$$

$$(e^{iH_0 t} V\Gamma(\Gamma^2(V)V)\Gamma(V))_{1,1}^{1PR}(p) + (V\Gamma(\Gamma(V)V)\Gamma^2(V)E^{-iH_0 t})_{1,1}^{1PR}(p)$$

To 4-th order of 1-particle average of wave operator give also contributions the 1PI diagrams, see Fig.5. The leading terms corresponding to these diagrams are of order t . We have

$$U_4^{1PI}(t, p) = it(V\Gamma(V\Gamma(V\Gamma(V))))_{1,1}^{1PI}(p) \quad (\text{B.11})$$

Above consideration proves that up to the t^2 -order the formula (4.2) is true.

References

- [1] N.N. Bogoliubov and D.V. Shirkov, *Introduction to the theory of quantum fields*, Nauka, 1973
- [2] S.S. Schweber, *An introduction to relativistic quantum field theory*, Row, Peterson and Co, N.Y. 1961.
- [3] K.O. Friedrichs, *Perturbation of Spectra in Hilbert Space*, AMS, Providence, 1965
- [4] L.D. Faddeev, Dokladi AN USSR, 152 (1963) 573
- [5] K. Hepp, *Theorie de la renormalisation*, Springer, 1969.
- [6] I.Ya. Aref'eva, Teor. Mat. Fis. 14 (1973) 3
- [7] J. Schwinger, *Field theory of unstable particles*, Ann. Phys. 9(1960)169-193
- [8] M.L. Goldberger and K.M. Watson, *Collision theory*, John Wiley & Sons, Inc, New York-London-Sydney, 1964
- [9] C. Cohen-Tannoudji, J. Dupont-Roc and G. Grinberg, *Atom-Photon Interactions, Basic Processes and Applications*, John Wiley & Sons, Inc., 1992
- [10] E.L. Feinberg, *A particle with non-equilibrium proper field*, in: Problems of theoretical physics, Memorial volume to Igor E. Tamm, Nauka, Moscow, 1972
- [11] V.A. Rubakov and M.E. Shaposhnikov, *Elektroweak baryon number non-conservation in the early Universe and in high-energy collisions*, Uspehi Fisicheskikh nauk, 166 (1996) 493-538
- [12] D.F. Walls and G.J. Milburn, *Quantum Optics*, Springer-Verlag, 1994.
- [13] W.H. Zurek, Phys. Rev. D26(1981)1516
- [14] I.V.Volovich, *Models of Quantum Computers and Decoherence Problem*, quant-ph/9902055; Proc. of the International Conference on Quantum Information, Meijo Univ. Nagoya, 4-8 Nov. 1997.
- [15] L. Accardi, I.Ya.Aref'eva and I.V.Volovich, *Non-Equilibrium Quantum Field Theory and Entangled Commutation Relations*, hep-th/9905035.
- [16] N.N. Bogoliubov, *Problems of dynamical theory in statistical physics*, Gostehizdat, 1946
- [17] K.O. Friedrichs, *On the perturbation of continuous spectra*, Comm. Pure Appl. Math. 1(1948)361-406.
- [18] L. van Hove, *Quantum mechanical perturbations giving rise to a transport equation*, Physica, 21(1955)517-540

- [19] I. Prigogine, *Non-equilibrium statistical mechanics* Pergamon, 1963,
- [20] D.N. Zubarev, *Non-equilibrium statistical thermodynamics*, Nauka, Moscow, 1971
- [21] E.M.Lifchiz and L.P.Pitaevskii, *Physical kinetics*, Nauka, Moscow, 1979
- [22] L. Accardi, S.V.Kozyrev, I.V. Volovich, *Dynamics of dissipative two-state systems in the stochastic approximation*, Phys. Rev. A 57 N. 3 (1997); quant-ph/9706021
- [23] L. Accardi, Y.G. Lu and I.V.Volovich, *Quantum theory and its stochastic limit*, Springer, (in press)
- [24] I.Ya. Aref'eva, Teor. Mat. Fis. 15 (1973) 207.